
| | |
|--------------------------|---|
| Product name: | ABCG2 Mouse Monoclonal Antibody |
| Cat number: | MABN80991 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Monoclonal |
| Concentration: | 1mg/ml |
| Host: | Mouse |
| Isotype: | Mouse IgG1 |
| Immunogen: | Purified recombinant fragment of human ABCG2 expressed in E. Coli. |
| Reactivity: | Human,Mouse,Rat,Rabbit,Monkey |
| Applications: | WB 1:500-1:2000,IHC 1:100-1:500,ICC 1:50-1:500,ELISA 1:5000-1:20000,FC 1:200-1:400 |
| Molecular Weight: | 72kDa |
| Purification: | Affinity Purification |
| Form: | Liquid |
| Buffer: | Purified antibody in PBS with 0.05% sodium azide. |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |
| Background: | <p>The membrane-associated protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Tissue specificity: Highly expressed in placenta. Low expression in small intestine, liver and colon.</p> |