

---

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | Abhd5 Rabbit Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN86650  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | A synthetic peptide of human Abhd5   |
| <b>Reactivity:</b>       | Human, Mouse   |
| <b>Applications:</b>     | WB 1:500-1:2000, IHC 1:500-1:2000, FC 1:100-1:200  |
| <b>Molecular Weight:</b> | Calculated MW:39 kDa; Observed MW:39 kDa   |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | The protein encoded by this gene belongs to a large family of proteins defined by an alpha/beta hydrolase fold, and contains three sequence motifs that correspond to a catalytic triad found in the esterase/lipase/thioesterase subfamily. It differs from other members of this subfamily in that its putative catalytic triad contains an asparagine instead of the serine residue. Mutations in this gene have been associated with Chanarin-Dorfman syndrome, a triglyceride storage disease with impaired long-chain fatty acid oxidation. [provided by RefSeq, Jul 2008] |