

---

<b>Product name:</b>	ADRM1 Rabbit Monoclonal Antibody
<b>Cat number:</b>	MABN85261
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A synthetic peptide of human ADRM1
<b>Reactivity:</b>	Human
<b>Applications:</b>	WB 1:500-1:1000,IHC 1:50-1:100,ICC 1:50-1:200,IP 1:10-1:20
<b>Molecular Weight:</b>	Calculated MW: 42 kDa; Observed MW: 42 kDa
<b>Purification:</b>	Affinity Purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Purified antibody in TBS with 0.05% sodium azide,0.05%BSA and 50% glycerol.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	<p>Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair. Within the complex, functions as a proteasomal ubiquitin receptor. Engages and activates 19S-associated deubiquitinases UCHL5 and PSMD14 during protein degradation. UCHL5 reversibly associate with the 19S regulatory particle whereas PSMD14 is an intrinsic subunit of the proteasome lid subcomplex.</p>