
Product name:	ARRDC3 Rabbit Polyclonal Antibody
Cat number:	ABN07167
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human ARRDC3. AA range:365-414
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight:	46kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	<p>This gene encodes a member of the arrestin family of proteins, which regulate G protein-mediated signaling. The encoded protein is thought to act as a regulator of breast cancer growth and progression by binding to a phosphorylated form of integrin beta4, a tumor-related antigen, targeting the integrin for internalization and degradation. [provided by RefSeq, Jul 2016],induction:By troglitazone and pioglitazone (selective PPARG agonists), by prostaglandin J2 (PGJ2) and by L165,041 (a PPARD ligand), by vitamin D3 and, to a lesser extent, by PMA (phorbol myristate acetate) in the promyelocytic leukemia HL-60 cells. No induction by retinoic acid, nor by clofibrate (a specific PPARA agonist).,similarity:Belongs to the arrestin family.,subcellular location:Associated with plasma membrane, as well as with endosomes and lysosomes during endocytosis.,subunit:Does not bind TXN (thioredoxin).,tissue specificity:Highly expressed in skeletal muscle, placenta, kidney, adrenal gland, lymph node, mammary gland, thyroid, and trachea. Very low levels in colon, thymus, spleen, small intestine, bladder and bone marrow. Strong expression in differentiated adipocytes compared to preadipocytes.,</p>