

---

<b>Product name:</b>	ATBF1 Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN07255
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	The antiserum was produced against synthesized peptide derived from human ZFH3. AA range:761-810
<b>Reactivity:</b>	Human,Mouse
<b>Applications:</b>	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	<p>This gene encodes a transcription factor with multiple homeodomains and zinc finger motifs, and regulates myogenic and neuronal differentiation. The encoded protein suppresses expression of the alpha-fetoprotein gene by binding to an AT-rich enhancer motif. The protein has also been shown to negatively regulate c-Myb, and transactivate the cell cycle inhibitor cyclin-dependent kinase inhibitor 1A (also known as p21CIP1). This gene is reported to function as a tumor suppressor in several cancers, and sequence variants of this gene are also associated with atrial fibrillation. Multiple transcript variants expressed from alternate promoters and encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2009],function:Transcriptional activator that binds to the AT-rich core sequence of the enhancer element of the AFP gene.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 22 C2H2-type zinc fingers.,similarity:Contains 4 homeobox DNA-binding domains.,subunit:Interacts with FNBP3.,</p>