

---

<b>Product name:</b>	ATG7 Rabbit Monoclonal Antibody
<b>Cat number:</b>	MABN21471
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG,Kappa
<b>Immunogen:</b>	A synthetic peptide of human ATG7
<b>Reactivity:</b>	Human,Mouse,Rat,
<b>Applications:</b>	WB 1:1000-1:5000,IHC 1:200-1:1000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
<b>Molecular Weight:</b>	Calculated MW:78kD;Observed MW:78kD
<b>Purification:</b>	Protein A
<b>Form:</b>	Liquid
<b>Buffer:</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	Cell localization:Cytoplasm.This gene encodes an E1-like activating enzyme that is essential for autophagy and cytoplasmic to vacuole transport. The encoded protein is also thought to modulate p53-dependent cell cycle pathways during prolonged metabolic stress. It has been associated with multiple functions, including axon membrane trafficking, axonal homeostasis, mitophagy, adipose differentiation, and hematopoietic stem cell maintenance. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015],