

<b>Product name:</b>	BAX
<b>Cat number:</b>	AB-10230
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	200 ug
<b>Clone:</b>	POLY
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthetic peptide mapping to the amino terminal domain of human Bax
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	WB: 1:1000 IP: 1:200 IHC(P): 1:200
<b>Purification:</b>	Aff. Pur
<b>Form:</b>	Liquid
<b>Buffer:</b>	20 mM Tris-HCl, pH 8.0
<b>Storage:</b>	Store this product at 4° C, do not freeze.
<b>Background:</b>	<p>Bcl-2 family of proteins is a key regulator of apoptosis that function to either inhibit or promote cell death. The over expression of members such as Bcl-2 and Bcl-xL inhibit the apoptotic process (1,2). The Bcl-2 family members are also characterized by dimerizing to further modulate apoptosis. Bag-1, for example, has been found to form a heterodimer with Bcl-2 resulting in the enhancement of the anti-apoptotic effect of Bcl-2 (3,4). Other anti-apoptotic Bcl-2 family members include A1, Bcl-xg, Bcl-xb, Mcl-1, BAR, BI-1 and Bcl-w (5). The pro-apoptotic family members include Bax, Bcl-xS, Bad, Bak, NBK, BID, Hrk, Bok, Bim, Noxa and Diva. Bax and Bak have been shown to play a critical role in cytochrome c release from mitochondria and thus initiate apoptosis (6). Bad plays a critical role in the Bax-mediated apoptosis pathway by dimerizing with Bcl-xL, causing the displacement of Bax. The displacement of Bax allows apoptosis to proceed (7). Bcl-xS, a shorter version of Bcl-xL (lacking amino acids 126-188), apparently utilizes a different pathway than Bax to induce cell death. Some research suggests that Bcl-xS uses a novel mechanism for regulating caspase or it may use an alternate cell death effector pathway (8,9).</p>

