
| | |
|--------------------------|---|
| Product name: | Bax Rabbit Monoclonal Antibody |
| Cat number: | MABN21477 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Monoclonal |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG,Kappa |
| Immunogen: | A synthetic peptide of human Bax |
| Reactivity: | Human,Mouse,Rat, |
| Applications: | WB 1:1000-1:5000,IHC 1:2000-1:5000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200 |
| Molecular Weight: | Calculated MW:21kD;Observed MW:21kD |
| Purification: | Protein A |
| Form: | Liquid |
| Buffer: | PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |
| Background: | Cell localization:Cytoplasm, Nuclear.The protein encoded by BAX (BCL2 associated X, apoptosis regulator) belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for BAX. |