



Product name:	C-Myc
Cat number:	MAB-94579
Size:	100µg
Clone:	MYC
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	Recombinant protein of human c-Myc
Reactivity:	Hu, Ms, Rt
Applications:	Western Blot: 1:500 - 1:2000 Immunohistochemistry: 1:50 - 1:100 Immunocytochemistry: 1:50 - 1:200 Immunofluorescence: 1:50 - 1:100 Immunoprecipitation: 1:20 - 1:50 Flow Cytometry: 1:20 - 1:50
Molecular Weight:	49kDa
Purification:	Aff. Pur.
Form:	Liquid
Buffer:	Store at -20°C. Avoid freeze / thaw cycles
Storage:	PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.
Background:	<p>The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene.</p>



Western blot analysis of MYC using anti-MYC antibody in 1) Jurkat cell lysate,(2) K562 cell lysate. The membrane was incubated with rabbit anti-MYC antigen affinity purified polyclonal antibody at 1 μ g/ml overnight at 4 $^{\circ}$ C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-Rabbit IgG IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an ECL West Pico Plus. A specific band was detected for MYC.



IHC analysis of MYC using anti-MYC antibody on paraffin-embedded human lung cancer.



Immunofluorescent analysis of HeLa cells, using c-Myc Antibody.