
| | |
|--------------------------|--|
| Product name: | Cadherin-22 Rabbit Polyclonal Antibody |
| Cat number: | ABN07831 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Polyclonal |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG |
| Immunogen: | The antiserum was produced against synthesized peptide derived from human CDH22. AA range:111-160 |
| Reactivity: | Human,Mouse,Rat |
| Applications: | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000 |
| Molecular Weight: | 84kDa |
| Purification: | Affinity purification |
| Form: | Liquid |
| Buffer: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |
| Background: | <p>This gene is a member of the cadherin superfamily. The gene product is composed of five cadherin repeat domains and a cytoplasmic tail similar to the highly conserved cytoplasmic region of classical cadherins. Expressed predominantly in the brain, this putative calcium-dependent cell adhesion protein may play an important role in morphogenesis and tissue formation in neural and non-neural cells during development and maintenance of the brain and neuroendocrine organs. [provided by RefSeq, Jul 2008],function:Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. PB-cadherins may have a role in the morphological organization of pituitary gland and brain tissues.,similarity:Contains 5 cadherin domains.,</p> |