
| | |
|--------------------------|---|
| Product name: | CD10 Rabbit Monoclonal Antibody |
| Cat number: | MABN87418 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Monoclonal |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG |
| Immunogen: | A synthetic peptide of human CD10 |
| Reactivity: | Human,Mouse,Rat |
| Applications: | WB 1:500-1:2000,IHC 1:200-1:1000,ICC/IF 1:20-1:50,IP 1:20-1:50 |
| Molecular Weight: | Calculated MW:86 kDa; Observed MW:100 kDa |
| Purification: | Affinity Purification |
| Form: | Liquid |
| Buffer: | Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |
| Background: | The protein encoded by this gene is a type II transmembrane glycoprotein and a common acute lymphocytic leukemia antigen that is an important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). The encoded protein is present on leukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. [provided by RefSeq, Aug 2017] |