

---

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | CD197 Mouse Monoclonal Antibody  |
| <b>Cat number:</b>       | MABN81972  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Mouse  |
| <b>Isotype:</b>          | Mouse IgG1   |
| <b>Immunogen:</b>        | Purified recombinant fragment of human CD197 (AA: extra mix) expressed in E. Coli.   |
| <b>Reactivity:</b>       | Human  |
| <b>Applications:</b>     | WB 1:500-1:2000,ELISA 1:5000-1:20000,FC 1:200-1:400  |
| <b>Molecular Weight:</b> | 42.9kDa  |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | <p>The protein encoded by this gene is a member of the G protein-coupled receptor family. This receptor was identified as a gene induced by the Epstein-Barr virus (EBV), and is thought to be a mediator of EBV effects on B lymphocytes. This receptor is expressed in various lymphoid tissues and activates B and T lymphocytes. It has been shown to control the migration of memory T cells to inflamed tissues, as well as stimulate dendritic cell maturation. The chemokine (C-C motif) ligand 19 (CCL19/ECL) has been reported to be a specific ligand of this receptor. Signals mediated by this receptor regulate T cell homeostasis in lymph nodes, and may also function in the activation and polarization of T cells, and in chronic inflammation pathogenesis. Alternative splicing of this gene results in multiple transcript variants.</p> |