

---

|                       |   |
|-----------------------|---|
| <b>Product name:</b>  | CD20 Mouse Monoclonal Antibody  |
| <b>Cat number:</b>    | MABN80615   |
| <b>Conjugate:</b>     | Unconjugated  |
| <b>Size:</b>          | 100µL   |
| <b>Clone:</b>         | Monoclonal  |
| <b>Concentration:</b> | 1mg/ml  |
| <b>Host:</b>          | Mouse   |
| <b>Isotype:</b>       | Mouse IgG   |
| <b>Immunogen:</b>     | Synthetic peptide corresponding to aa (EPANPSEKNSPSTQY) of human CD20, conjugated to KLH.   |
| <b>Reactivity:</b>    | Human   |
| <b>Applications:</b>  | IHC 1:200-1:1000, ICC 1:200-1:1000, ELISA 1:5000-1:20000  |
| <b>Purification:</b>  | Affinity Purification   |
| <b>Form:</b>          | Liquid  |
| <b>Buffer:</b>        | Ascitic fluid containing 0.03% sodium azide.  |
| <b>Storage:</b>       | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>    | CD20 (MS4A1, membrane-spanning 4-domains, subfamily A, member 1) is a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. The CD20 antigen is present on human pre B lymphocytes and on B lymphocytes at all stages of maturation, except on plasma cells. Low level expression of the CD20 antigen has been detected on normal T lymphocytes. The CD20 molecule is involved in regulation of B cell differentiation, presumably via its reported function as a Ca <sup>++</sup> channel subunit. And it is known to accelerate the G0 to G1 progression induced by IGF-1. |