

---

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | CD5 Mouse Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN82532   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Mouse   |
| <b>Isotype:</b>          | Mouse IgG1  |
| <b>Immunogen:</b>        | Purified recombinant fragment of human CD5 (AA: 403-495) expressed in E. Coli.  |
| <b>Reactivity:</b>       | Human, Mouse  |
| <b>Applications:</b>     | WB 1:500-1:2000,ELISA 1:5000-1:20000,FC 1:200-1:400   |
| <b>Molecular Weight:</b> | 54.6kDa   |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | This gene encodes a member of the scavenger receptor cysteine-rich (SRCR) superfamily. Members of this family are secreted or membrane-anchored proteins mainly found in cells associated with the immune system. This protein is a type-I transmembrane glycoprotein found on the surface of thymocytes, T lymphocytes and a subset of B lymphocytes. The encoded protein contains three SRCR domains and may act as a receptor to regulate T-cell proliferation. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Oct 2016] |