

---

|                       |  |
|-----------------------|--|
| <b>Product name:</b>  | Chemokine Receptor D6 Rabbit Polyclonal Antibody   |
| <b>Cat number:</b>    | ABN08753   |
| <b>Conjugate:</b>     | Unconjugated   |
| <b>Size:</b>          | 100µL  |
| <b>Clone:</b>         | Polyclonal   |
| <b>Concentration:</b> | 1mg/ml   |
| <b>Host:</b>          | Rabbit   |
| <b>Isotype:</b>       | IgG  |
| <b>Immunogen:</b>     | The antiserum was produced against synthesized peptide derived from human CCBP2. AA range:335-384  |
| <b>Reactivity:</b>    | Human,Rat,Mouse  |
| <b>Applications:</b>  | ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000  |
| <b>Purification:</b>  | Affinity purification  |
| <b>Form:</b>          | Liquid   |
| <b>Buffer:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.   |
| <b>Storage:</b>       | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>    | <p>This gene encodes a beta chemokine receptor, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptor-mediated signal transduction are critical for the recruitment of effector immune cells to the inflammation site. This gene is expressed in a range of tissues and hemopoietic cells. The expression of this receptor in lymphatic endothelial cells and overexpression in vascular tumors suggested its function in chemokine-driven recirculation of leukocytes and possible chemokine effects on the development and growth of vascular tumors. This receptor appears to bind the majority of beta-chemokine family members; however, its specific function remains unknown. This gene is mapped to chromosome 3p21.3, a region that includes a cluster of chemokine receptor genes. [provided by RefSeq, Jul 2008],function:Receptor for C-C type chemokines including SCYA2/MCP-1, SCY3/MIP-1-alpha, SCYA5/RANTES and SCYA7/MCP-3.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed primarily in placenta and fetal liver, and found at very low levels in the lung and lymph node. Found in endothelial cells lining afferent lymphatics in dermis and lymph nodes. Also found in lymph nodes subcapsular and medullary sinuses, tonsillar lymphatic sinuses and lymphatics in mucosa and submucosa of small and large intestine and appendix. Also found in some malignant vascular tumors.,</p> |