

Product name:	CLECSF6 Rabbit Polyclonal Antibody
Cat number:	ABN09035
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from the C-terminal region of human CLEC4A. AA range:161-210
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight:	28kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. The encoded type 2 transmembrane protein may play a role in inflammatory and immune response. Multiple transcript variants encoding distinct isoforms have been identified for this gene. This gene is closely linked to other CTL/CTLD superfamily members on chromosome 12p13 in the natural killer gene complex region. [provided by RefSeq, Jul 2008],domain:Contains 1 copy of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,function:May be involved in regulating immune reactivity. May play a role in modulating dendritic cells (DC) differentiation and/or maturation. May be involved via its ITIM motif (immunoreceptor tyrosine-based inhibitory motifs) in the inhibition of B-cell-receptor-mediated calcium mobilization and protein tyrosine phosphorylation.,induction:Agonists of neutrophils GM-CSF, IL-3, IL-4 and IL-13 up-regulated mRNA expression of isoform 2.,online information:DCIR,similarity:Contains 1 C-type lectin domain.,subunit:May interact with PTPN6 via its ITIM motif.,tissue specificity:Expressed in dendritic cells, myeloid cells, B-cells and HL-60 cells (at protein level). TNF alpha, IL-1 alpha, and LPS, down-regulated expression at the surface of neutrophils (at protein level). Expressed preferentially in hematopoietic tissues. Expressed in peripheral blood leukocytes, neutrophils, moderate quantities in spleen, lymph node, and bone marrow, and at very low levels in thymus. Expressed in Ag-presenting cells (DC, monocytes, macrophages and B-cells), as well as on granulocytes. Expression was decreased in DC by signals inducing its maturation (e.g. CD40 ligand, LPS, and TNF alpha),.