

Product name:	CLIC3 Rabbit Polyclonal Antibody
Cat number:	ABN09038
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human CLIC3. AA range:21-70
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:20000-1:40000
Molecular Weight:	27kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

chloride intracellular channel 3 (CLIC3) Homo sapiens Chloride channels are a diverse group of proteins that regulate fundamental cellular processes including stabilization of cell membrane potential, transepithelial transport, maintenance of intracellular pH, and regulation of cell volume. Chloride intracellular channel 3 is a member of the p64 family and is predominantly localized in the nucleus and stimulates chloride ion channel activity. In addition, this protein may participate in cellular growth control, based on its association with ERK7, a member of the MAP kinase family. [provided by RefSeq, Jul 2008], domain: Members of this family may change from a globular, soluble state to a state where the N-terminal domain is inserted into the membrane and functions as chloride channel. A conformation change of the N-terminal domain is thought to expose hydrophobic surfaces that trigger membrane insertion., function: Can insert into membranes and form chloride ion channels. May participate in cellular growth control., similarity: Belongs to the chloride channel CLIC family., similarity: Contains 1 GST C-terminal domain., similarity: Contains 1 GST N-terminal domain., subcellular location: Predominantly nuclear. Some protein was found in the cytoplasm. Exists both as soluble cytoplasmic protein and as membrane protein with probably a single transmembrane domain., subunit: Associated with the C-terminal of ERK7., tissue specificity: Detected in placenta (at protein level). Widely expressed. High expression is found in placenta followed by lung and heart. Low expression in skeletal muscle, kidney and pancreas.,