

Product name:	CPN cat Rabbit Polyclonal Antibody
Cat number:	ABN09320
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human CPN1. AA range:409-458
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight:	total 52kDa,Cleaved 48kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

Carboxypeptidase N is a plasma metallo-protease that cleaves basic amino acids from the C terminal of peptides and proteins. The enzyme is important in the regulation of peptides like kinins and anaphylatoxins, and has also been known as kininase-1 and anaphylatoxin inactivator. This enzyme is a tetramer comprised of two identical regulatory subunits and two identical catalytic subunits; this gene encodes the catalytic subunit. Mutations in this gene can be associated with angioedema or chronic urticaria resulting from carboxypeptidase N deficiency. [provided by RefSeq, Jul 2008],catalytic activity:Release of a C-terminal basic amino acid, preferentially lysine.,cofactor:Binds 1 zinc ion per subunit.,disease:Defects in CPN1 are the cause of carboxypeptidase N deficiency [MIM:212070]. Patients affected present some combination of angioedema or chronic urticaria, as well as hay fever or asthma, and have also slightly depressed serum carboxy peptidase N, suggestive of autosomal recessive inheritance of this disorder.,function:Protects the body from potent vasoactive and inflammatory peptides containing C-terminal Arg or Lys (such as kinins or anaphylatoxins) which are released into the circulation.,similarity:Belongs to the peptidase M14 family.,subunit:Tetramer of two catalytic chains and two glycosylated inactive chains.,tissue specificity:Synthesized in the liver and secreted in plasma.,