

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | CXCR3 Rabbit Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN83985  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | A synthesized peptide derived from human CXCR3   |
| <b>Reactivity:</b>       | Human  |
| <b>Applications:</b>     | WB 1:1000-1:2000,ICC 1:50-1:200,IP 1:20-1:50   |
| <b>Molecular Weight:</b> | 41 kDa   |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide,0.05% BSA and 50% glycerol.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | Receptor for the C-X-C chemokine CXCL9, CXCL10 and CXCL11 and mediates the proliferation, survival and angiogenic activity of human mesangial cells (HMC) through a heterotrimeric G-protein signaling pathway.Binds to CCL21. Probably promotes cell chemotaxis response. Upon activation by PF4, induces activated T-lymphocytes migration mediated via downstream Ras/extracellular signal-regulated kinase (ERK) signaling |