

Product name:	CYP17A1 Rabbit Polyclonal Antibody
Cat number:	ABN09627
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human Cytochrome P450 17A1. AA range:221-270
Reactivity:	Human
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
Molecular Weight:	50kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

cytochrome P450 family 17 subfamily A member 1(CYP17A1) Homo sapiens This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum. It has both 17alpha-hydroxylase and 17,20-lyase activities and is a key enzyme in the steroidogenic pathway that produces progestins, mineralocorticoids, glucocorticoids, androgens, and estrogens. Mutations in this gene are associated with isolated steroid-17 alpha-hydroxylase deficiency, 17-alpha-hydroxylase/17,20-lyase deficiency, pseudohermaphroditism, and adrenal hyperplasia. [provided by RefSeq, Jul 2008],catalytic activity:A steroid + AH(2) + O(2) = a 17-alpha-hydroxysteroid + A + H(2)O.,cofactor:Heme group.,disease:Defects in CYP17A1 are the cause of adrenal hyperplasia type 5 (AH5) [MIM:202110]. AH5 is a form of congenital adrenal hyperplasia, a common recessive disease due to defective synthesis of cortisol. Congenital adrenal hyperplasia is characterized by androgen excess leading to ambiguous genitalia in affected females, rapid somatic growth during childhood in both sexes with premature closure of the epiphyses and short adult stature. Four clinical types: "salt wasting" (SW, the most severe type), "simple virilizing" (SV, less severely affected patients), with normal aldosterone biosynthesis, "non-classic form" or late onset (NC or LOAH), and "cryptic" (asymptomatic).,enzyme regulation:Regulated predominantly by intracellular cAMP levels.,function:Conversion of pregnenolone and progesterone to their 17-alpha-hydroxylated products and subsequently to dehydroepiandrosterone (DHEA) and androstenedione. Catalyzes both the 17-alpha-hydroxylation and the 17,20-lyase reaction. Involved in sexual development during fetal life and at puberty.,online information:The Singapore human mutation and polymorphism database,pathway:Lipid metabolism; steroid biosynthesis.,PTM:Phosphorylation is necessary for 17,20-lyase, but not for 17-alpha-hydroxylase activity.,similarity:Belongs to the cytochrome P450 family.,