

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | Cystatin C Mouse Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN84979  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Mouse  |
| <b>Isotype:</b>          | Mouse IgG1   |
| <b>Immunogen:</b>        | Purified recombinant protein expressed in E.coli.  |
| <b>Reactivity:</b>       | Human  |
| <b>Applications:</b>     | WB 1:500-1:1000,ELISA 1:5000-1:20000   |
| <b>Molecular Weight:</b> | Calculated MW: 16 kDa; Observed MW: 16 kDa   |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide,0.5%BSA and 50% glycerol.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | Cystatin C is a 14 kDa member of the Cystatin superfamily of cysteine protease inhibitors. Most cell types secrete Cystatin C. Cystatin C inhibits cathepsins, and thereby may function as a tumor suppressor by inhibiting cathepsin mediated tumor cell invasion. In addition, this tumor suppressor function can also be attributed to Cystatin C's ability to antagonize TGF-β1 signaling. |