
Product name:	DHA Kinase Rabbit Polyclonal Antibody
Cat number:	ABN09958
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human DAK. AA range:91-140
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:200-1:1000,IHC 1:100-1:500,ICC/IF 1:20-1:50,ELISA 1:10000-1:20000
Molecular Weight:	59kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	<p>This gene is a member of the family of dihydroxyacetone kinases, which have a protein structure distinct from other kinases. The product of this gene phosphorylates dihydroxyacetone, and also catalyzes the formation of riboflavin 4',5'-phosphate (aka cyclin FMN) from FAD. Several alternatively spliced transcript variants have been identified, but the full-length nature of only one has been determined. [provided by RefSeq, Jul 2008],catalytic activity:ATP + glycerone = ADP + glycerone phosphate.,catalytic activity:FAD = AMP + riboflavin cyclic-4',5'-phosphate.,cofactor:Magnesium.,cofactor:Manganese or cobalt; for FAD-AMP lyase activity.,enzyme regulation:Each activity is inhibited by the substrate(s) of the other.,function:Catalyzes both the phosphorylation of dihydroxyacetone and the splitting of ribonucleoside diphosphate-X compounds among which FAD is the best substrate.,similarity:Belongs to the dihydroxyacetone kinase (DAK) family.,similarity:Contains 1 DAK1 (dihydroxyacetone kinase subunit 1) domain.,similarity:Contains 1 DAK2 (dihydroxyacetone kinase subunit 2) domain.,subunit:Homodimer.,</p>