

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | EAAT2 Rabbit Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN86785  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | A synthetic peptide of mouse EAAT2   |
| <b>Reactivity:</b>       | Mouse,Rat  |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:200-1:2000,IP 1:20-1:50  |
| <b>Molecular Weight:</b> | Calculated MW:62 kDa; Observed MW:65 kDa   |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed:7698742, PubMed:7557442, PubMed:9373176). Functions as a symporter that transports one amino acid molecule together with two or three Na <sup>+</sup> ions and one proton, in parallel with the counter-transport of one K <sup>+</sup> ion. Mediates Cl <sup>-</sup> flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na <sup>+</sup> symport (By similarity). Essential for the rapid removal of released glutamate from the synaptic cleft, and for terminating the postsynaptic action of glutamate (PubMed:9180080). |