

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | EIF4B Mouse Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN81132   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Mouse   |
| <b>Isotype:</b>          | Mouse IgG2b   |
| <b>Immunogen:</b>        | Purified recombinant fragment of human EIF4B expressed in E. Coli.  |
| <b>Reactivity:</b>       | Human, Mouse  |
| <b>Applications:</b>     | WB 1:500-1:2000, IHC 1:200-1:1000, ICC 1:200-1:1000, ELISA 1:5000-1:20000, FC 1:200-1:400   |
| <b>Molecular Weight:</b> | 80kDa   |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | The eukaryotic translation initiation factor 4B (eIF4B) plays a critical role in recruiting the 40S ribosomal subunit to the mRNA. It functions in close association with eIF4F and eIF4A. It binds near the 5'-terminal cap of mRNA in the presence of eIF4F and ATP. It promotes the ATPase activity and the ATP-dependent RNA unwinding activity of both eIF4A and eIF4F |