

<b>Product name:</b>	Eps15 (phospho Tyr849) Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN04623
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthesized phospho-peptide around the phosphorylation site of human Eps15 (phospho Tyr849)
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Applications:</b>	WB 1:500-1:2000,ELISA 1:5000-1:20000
<b>Molecular Weight:</b>	100kDa
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**Background:**

This gene encodes a protein that is part of the EGFR pathway. The protein is present at clathrin-coated pits and is involved in receptor-mediated endocytosis of EGF. Notably, this gene is rearranged with the HRX/ALL/MLL gene in acute myelogenous leukemias. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, May 2009],disease:A chromosomal aberration involving EPS15 is found in acute leukemias. Translocation t(1;11)(p32;q23) with MLL/HRX. The result is a rogue activator protein.,domain:The EH domain interacts with Asn-Pro-Phe (NPF) motifs of target proteins.,function:Involved in cell growth regulation. May be involved in the regulation of mitogenic signals and control of cell proliferation. Involved in the internalization of ligand-inducible receptors of the receptor tyrosine kinase (RTK) type, in particular EGFR. Plays a role in the assembly of clathrin-coated pits.,PTM:Phosphorylation on Tyr-849 is involved in the internalization of EGFR. Not required for membrane translocation after EGF treatment or for targeting to coated pits, but essential for a subsequent step in EGFR endocytosis (By similarity). Phosphorylated on serine upon DNA damage, probably by ATM or ATR.,similarity:Contains 2 EF-hand domains.,similarity:Contains 2 UIM (ubiquitin-interacting motif) repeats.,similarity:Contains 3 EH domains.,subcellular location:Colocalizes with HGS on bilayered clathrin coats on endosomes.,subcellular location:Recruited to the plasma membrane upon EGFR activation and localizes to coated pits.,subunit:Interacts with HGS; the interaction bridges the interaction of STAM or STAM2 with EPS15. Isoform 2 interacts with HGS and AP2A2. Part of a complex at least composed of EPS15, HGS, and either STAM1 or STAM2. Binds AP2A2 and AP2B1. Binds STON2 and EPN1. Interacts (via its SH3-binding sites) with CRK. Interacts with SH3BP4/TTP.,tissue specificity:Ubiquitously expressed.,