

---

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | Estrogen Receptor alpha Rabbit Monoclonal antibody  |
| <b>Cat number:</b>       | MABN01960   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Rabbit  |
| <b>Isotype:</b>          | IgG   |
| <b>Immunogen:</b>        | A synthetic peptide of human Estrogen Receptor alpha  |
| <b>Reactivity:</b>       | Human   |
| <b>Applications:</b>     | WB 1:500-1:1000,IHC 1:50-1:100,ICC/IF 1:50-1:200,IP 1:20-1:50,ChIP 1:20   |
| <b>Molecular Weight:</b> | Calculated MW: 66 kDa; Observed MW: 66 kDa  |
| <b>Purification:</b>     | Affinity Purified   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | ER (estrogen receptor 1) a member of the steroid receptor superfamily, contains highly conserved DNA binding (DBD) and ligand binding domains (LBD). Through its estrogen-independent and estrogen-dependent activation domains (AF-1 and AF-2, respectively), ER regulates transcription by recruiting coactivator proteins and interacting with general transcriptional machinery. Phosphorylation provides an important mechanism to regulate ER activity. ER is phosphorylated on multiple sites. |