

---

<b>Product name:</b>	FAST Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN10840
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	The antiserum was produced against synthesized peptide derived from human FAST. AA range:361-410
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Applications:</b>	WB 1:500-1:2000,ELISA 1:10000-1:20000
<b>Molecular Weight:</b>	61kDa
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	<p>The protein encoded by this gene is a member of the serine/threonine protein kinase family. This kinase was shown to be activated rapidly during Fas-mediated apoptosis in Jurkat cells. In response to Fas receptor ligation, it phosphorylates TIA1, an apoptosis-promoting nuclear RNA-binding protein. The encoded protein is a strong inducer of lymphocyte apoptosis. Two transcript variants encoding different isoforms have been found for this gene. Other variants exist, but their full-length natures have not yet been determined. [provided by RefSeq, Jul 2008],catalytic activity:ATP + [Fas-activated serine/threonine protein] = ADP + [Fas-activated serine/threonine phosphoprotein].,function:Fas-mediated apoptosis is characterized by FASTK-dephosphorylation and TIA-1-phosphorylation. Both are activated and FASTK-mediated TIA-1 activation play a key role in apoptosis.,PTM:Phosphorylated on serine/threonine.,similarity:Belongs to the FAST protein kinase family.,similarity:Contains 1 RAP domain.,subunit:Interacts with TIA-1.,tissue specificity:Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.,</p>