

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | Fractalkine Receptor Rabbit Polyclonal Antibody  |
| <b>Cat number:</b>       | ABN00484   |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Polyclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | Synthesized peptide derived from Fractalkine Receptor . at AA range: 120-200   |
| <b>Reactivity:</b>       | Human,Mouse,Rat  |
| <b>Applications:</b>     | WB 1:500-1:1000,IHC 1:50-1:100,ELISA 1:5000-1:20000  |
| <b>Molecular Weight:</b> | Calculated MW: 40 kDa; Observed MW: 40 kDa   |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | Receptor for the C-X3-C chemokine fractalkine (CX3CL1) present on many early leukocyte cells; CX3CR1-CX3CL1 signaling exerts distinct functions in different tissue compartments, such as immune response, inflammation, cell adhesion and chemotaxis. |