
| | |
|--------------------------|--|
| Product name: | Galectin-9 Rabbit Polyclonal Antibody |
| Cat number: | ABN11278 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Polyclonal |
| Concentration: | 1mg/ml |
| Host: | Rabbit |
| Isotype: | IgG |
| Immunogen: | The antiserum was produced against synthesized peptide derived from human LEG9. AA range:51-100 |
| Reactivity: | Human,Mouse,Rat |
| Applications: | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000 |
| Molecular Weight: | 40kDa |
| Purification: | Affinity purification |
| Form: | Liquid |
| Buffer: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |
| Background: | <p>The galectins are a family of beta-galactoside-binding proteins implicated in modulating cell-cell and cell-matrix interactions. The protein encoded by this gene is an S-type lectin. It is overexpressed in Hodgkin's disease tissue and might participate in the interaction between the H&RS cells with their surrounding cells and might thus play a role in the pathogenesis of this disease and/or its associated immunodeficiency. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2008],alternative products:Additional isoforms seem to exist,domain:Contains two homologous but distinct carbohydrate-binding domains.,function:Binds galactosides. Has high affinity for the Forssman pentasaccharide. May play a role in thymocyte-epithelial interactions relevant to the biology of the thymus. Inhibits cell proliferation. The isoform Short acts as an eosinophil chemoattractant. Is a ligand for HAVCR2/TIM3. Induces T-helper type 1 lymphocyte (Th1) death.,online information:Galectin-9,similarity:Contains 2 galectin domains.,subcellular location:May also be secreted by a non-classical secretory pathway.,subunit:Monomer.,tissue specificity:Peripheral blood leukocytes and lymphatic tissues. Overexpressed in Hodgkin's disease tissue.,</p> |