

---

<b>Product name:</b>	Glycerate Kinase Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN11511
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	The antiserum was produced against synthesized peptide derived from human GLCTK. AA range:101-150
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Applications:</b>	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
<b>Molecular Weight:</b>	60kDa
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	<p>This locus encodes a member of the glycerate kinase type-2 family. The encoded enzyme catalyzes the phosphorylation of (R)-glycerate and may be involved in serine degradation and fructose metabolism. Decreased activity of the encoded enzyme may be associated with the disease D-glyceric aciduria. Alternatively spliced transcript variants have been described. [provided by RefSeq, Jan 2009],catalytic activity:ATP + (R)-glycerate = ADP + 3-phospho-(R)-glycerate.,disease:Defects in GLYCTK are the cause of D-glyceric acidemia [MIM:220120]. It is characterized by nonketotic hyperglycinemia with the excretion of D-glyceric acid in the urine and the presence of this substance in the serum.,similarity:Belongs to the glycerate kinase type-2 family.,tissue specificity:Widely expressed.,</p>