

---

<b>Product name:</b>	Histone H2B (yeast) Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN00044
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Reactivity:</b>	Yeast
<b>Applications:</b>	WB 1:500-1:1000
<b>Molecular Weight:</b>	Calculated MW: 14 kDa; Observed MW: 14 kDa
<b>Purification:</b>	Affinity Chromatography
<b>Form:</b>	Liquid
<b>Buffer:</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.