

---

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | HMGB1 Rabbit Monoclonal Antibody  |
| <b>Cat number:</b>       | MABN87283   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Rabbit  |
| <b>Isotype:</b>          | IgG   |
| <b>Immunogen:</b>        | A synthetic peptide of human HMGB1  |
| <b>Reactivity:</b>       | Human,Mouse,Rat   |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:200,FC 1:20-1:100   |
| <b>Molecular Weight:</b> | Calculated MW:25 kDa; Observed MW:25 kDa  |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | <p>This gene encodes a protein that belongs to the High Mobility Group-box superfamily. The encoded non-histone, nuclear DNA-binding protein regulates transcription, and is involved in organization of DNA. This protein plays a role in several cellular processes, including inflammation, cell differentiation and tumor cell migration. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Sep 2015]</p> |