

---

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | HTR3A Mouse Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN81929   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Mouse   |
| <b>Isotype:</b>          | Mouse IgG2a   |
| <b>Immunogen:</b>        | Purified recombinant fragment of human HTR3A (AA: extra 24-157) expressed in E. Coli.   |
| <b>Reactivity:</b>       | Human   |
| <b>Applications:</b>     | IHC 1:200-1:1000, ICC 1:100-1:500, ELISA 1:5000-1:20000, FC 1:200-1:400   |
| <b>Molecular Weight:</b> | 55.3kDa   |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | The product of this gene belongs to the ligand-gated ion channel receptor superfamily. This gene encodes subunit A of the type 3 receptor for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. This receptor causes fast, depolarizing responses in neurons after activation. It appears that the heteromeric combination of A and B subunits is necessary to provide the full functional features of this receptor, since either subunit alone results in receptors with very low conductance and response amplitude. Alternatively spliced transcript variants encoding different isoforms have been identified. |