
Product name:	Interferon Receptor alpha (15I18) Rabbit Monoclonal Antibody
Cat number:	MABN12685
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human Interferon alpha/beta receptor 1
Reactivity:	Human
Applications:	WB 1:1000-1:5000
Molecular Weight:	64kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	Interferon Receptor alpha (IFN-α R1) is a class II cytokine receptor which belongs to type I human interferons (IFNs) family. IFNs plays a role in antiviral, antiproliferative, immunomodulatory, antitumor, and antiparasitic activities by inducing transcription of IFN-stimulated genes (ISGs) through activation of the Jak-STAT pathway. Component of the receptor for type I interferons, including interferons alpha, IFNB1 and IFNW1 (PubMed:2153461, PubMed:7665574, PubMed:10049744, PubMed:14532120, PubMed:15337770, PubMed:21854986). Functions in general as heterodimer with IFNAR2 (PubMed:7665574, PubMed:10049744, PubMed:21854986). Type I interferon binding activates the JAK-STAT signaling cascade, and triggers tyrosine phosphorylation of a number of proteins including JAKs, TYK2, STAT proteins and the IFNR alpha- and beta-subunits themselves (PubMed:7665574, PubMed:21854986, PubMed:32972995). Can form an active IFNB1 receptor by itself and activate a signaling cascade that does not involve activation of the JAK-STAT pathway (By similarity).