




Product name:	Ki-67
Cat number:	AB-90972
Conjugate:	Unconjugated
Size:	100 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	Recombinant human construct containing amino acids 1,111-1,490 expressed in and purified from E. coli.
Reactivity:	Hu, Ms, Rt
Applications:	Western Blot : 1:2,000-1:5,000 Immunohistochemistry (paraffin, formalin, frozen Tissues): 1:500 Immunofluorescence/Immunocytochemistry: 1:1,000-1:2,500
Molecular Weight:	345-395kDa
Purification:	serum
Form:	Liquid
Buffer:	Supplied as an aliquot of serum plus 5mM sodium azide.
Storage:	Storage for short term at 4°C recommended, for longer term at -20°C, minimize freeze/thaw cycles
Background:	<p>The Ki-67 proteins were first discovered in an attempt to generate cancer specific monoclonal antibodies. A monoclonal antibody which bound to structures in the nuclei of dividing but not quiescent cells was produced and shown to bind two very large proteins of molecular weight 345kDa and 395kDa. The two proteins were derived from alternate transcripts of a single gene. The presence of Ki-67 proteins, detected with an appropriate antibody, is an indicator of cell proliferation and the level of Ki-67 expression is one of the most reliable biomarkers of proliferative status of cancer cells. The RPCA-Ki-67 antibody was raised against a recombinant construct containing amino acids 1,111-1,490 of human Ki-67 isotype 1. The antibody can be used to identify dividing cells in rat and mouse brain and also works on paraffin sections of human tissues, where it is useful to identify cancer cells. Mouse select image at left for larger view.</p>




Immunofluorescent analysis of HeLa cells stained with rabbit Anti-Ki-67 dilution 1:2,500 in red, and mouse monoclonal antibody to Fibrillarin, dilution 1:2,000, in green. The blue is DAPI staining of nuclear DNA.



High magnification confocal image of adult mouse hippocampus dentate region stained with Anti-Ki-67 Polyclonal 1:2,000 - in green. Blue is Hoechst dye staining of DNA. Top left is DNA, top right FOX3/NeuN, bottom left Ki-67 and bottom right all three merged. Dividing cells are very rare in adult animals, but one can be seen in the center of the image. Chromosomes can be seen in blue and their Ki-67 coating can be seen in red. The dividing cell is FOX3/NeuN negative and so is presumably a glial cell.



Mouse NIH-3T3 cells stained with Anti-Ki-67 1:2,500 in red and mouse mAb to β -tubulin, 1:1,000 in green. The Ki-67 strongly stains the nuclei of dividing cells, but not quiescent cells



A section of human breast tissue including both normal and cancer cells. The cancer cells divide rapidly and heavily express Ki-67 and so stain strongly with the Anti-Ki-67 antibody.