

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | KMT2C Mouse Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN82259   |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Monoclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Mouse   |
| <b>Isotype:</b>          | Mouse IgG1  |
| <b>Immunogen:</b>        | Purified recombinant fragment of human KMT2C (AA: 1-205) expressed in E. Coli.  |
| <b>Reactivity:</b>       | Human   |
| <b>Applications:</b>     | ELISA 1:5000-1:20000,FC 1:200-1:400   |
| <b>Molecular Weight:</b> | 541.4kDa  |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | This gene is a member of the myeloid/lymphoid or mixed-lineage leukemia (MLL) family and encodes a nuclear protein with an AT hook DNA-binding domain, a DHHC-type zinc finger, six PHD-type zinc fingers, a SET domain, a post-SET domain and a RING-type zinc finger. This protein is a member of the ASC-2/NCOA6 complex (ASCOM), which possesses histone methylation activity and is involved in transcriptional coactivation. [provided by RefSeq, Jul 2008] |