

Product name:	Kv4.2 Rabbit Polyclonal Antibody
Cat number:	ABN13169
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human Kv4.2/KCND2. AA range:581-630
Reactivity:	Human,Mouse,Rat
Applications:	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. Four sequence-related potassium channel genes - shaker, shaw, shab, and shal - have been identified in *Drosophila*, and each has been shown to have human homolog(s). This gene encodes a member of the potassium channel, voltage-gated, shal-related subfamily, members of which form voltage-activated A-type potassium ion channels and are prominent in the repolarization phase of the action potential. This member mediates a rapidly inactivating, A-type outward potassium current which is not under the control of the N terminus as is domain: The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position., function: Pore-forming (alpha) subunit of voltage-gated rapidly inactivating A-type potassium channels. May contribute to I(To) current in heart and I(Sa) current in neurons. Channel properties are modulated by interactions with other alpha subunits and with regulatory subunits., PTM: Phosphorylated on serine and threonine residues., similarity: Belongs to the potassium channel family. D (Shal) subfamily., subcellular location: Detected in dendrites in cultured hippocampal neurons. Association with KCNIP2 probably enhances cell surface expression., subunit: Homotetramer or heterotetramer with KCND1 and/or KCND3. Interacts with DPP6, DLG4 and FREQ (By similarity). Interacts with DLG1. Associates with the regulatory subunits KCNIP1, KCNIP2, KCNIP3 and KCNIP4. Probably part of a complex consisting of KCNIP1, KCNIP2 isoform 3 and KCND2. The KCND2-KCNIP2 channel complex contains four KCND2 and four KCNIP2 subunits. Interacts with FLNA, FLNC and DPP10., tissue specificity: Highly expressed throughout the brain. Expression is very low or absent in other tissues.,