

---

<b>Product name:</b>	LRBA Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN13401
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Synthesized peptide derived from human protein . at AA range: 1970-2050
<b>Reactivity:</b>	Human,Mouse
<b>Applications:</b>	IHC 1:50-1:300,ICC/IF 1:50-1:200
<b>Molecular Weight:</b>	314kDa
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	<p>The protein encoded by this gene is a member of the WDL-BEACH-WD (WBW) gene family. Its expression is induced in B cells and macrophages by bacterial lipopolysaccharides (LPS). The encoded protein associates with protein kinase A and may be involved in leading intracellular vesicles to activated receptor complexes, which aids in the secretion and/or membrane deposition of immune effector molecules. Defects in this gene are associated with the disorder common variable immunodeficiency-8 with autoimmunity. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012],caution:Was originally (PubMed:1505956) said to be similar to yeast CDC4, but that similarity is very limited.,induction:By lipopolysaccharide (LPS).,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 BEACH domain.,similarity:Contains 6 WD repeats.,</p>