
Product name:	Lysophospholipase 1 (18C9) Rabbit Monoclonal Antibody
Cat number:	MABN04965
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human Lysophospholipase 1/LPL-I
Reactivity:	Human, Mouse, Rat
Applications:	WB 1:500-1:2000
Molecular Weight:	25kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	Hydrolyzes fatty acids from S-acylated cysteine residues in proteins such as trimeric G alpha proteins or HRAS. Has depalmitoylating activity and also low lysophospholipase activity. Acts as a acyl-protein thioesterase (PubMed:19439193, PubMed:20418879). Hydrolyzes fatty acids from S-acylated cysteine residues in proteins such as trimeric G alpha proteins or HRAS (PubMed:20418879). Has depalmitoylating activity toward KCNMA1 (PubMed:22399288). Could also depalmitoylate ADRB2 (PubMed:27481942). Acts as a lysophospholipase and hydrolyzes lysophosphatidylcholine (lyso-PC) (PubMed:19439193). Also hydrolyzes lysophosphatidylethanolamine (lyso-PE), lysophosphatidylinositol (lyso- PI) and lysophosphatidylserine (lyso-PS) (By similarity). Has much higher thioesterase activity than lysophospholipase activity (PubMed:19439193). Contributes to the production of lysophosphatidic acid (LPA) during blood coagulation by recognizing and cleaving plasma phospholipids to generate lysophospholipids which in turn act as substrates for ENPP2 to produce LPA (PubMed:21393252).