

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | LZK Rabbit Polyclonal Antibody  |
| <b>Cat number:</b>       | ABN13525  |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Polyclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Rabbit  |
| <b>Isotype:</b>          | IgG   |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human M3K13. AA range:151-200 |
| <b>Reactivity:</b>       | Human,Mouse,Rat   |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000                           |
| <b>Molecular Weight:</b> | 108kDa  |
| <b>Purification:</b>     | Affinity purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.                |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.      |

**Background:**

The protein encoded by this gene is a member of serine/threonine protein kinase family. This kinase contains a dual leucine-zipper motif, and has been shown to form dimers/oligomers through its leucine-zipper motif. This kinase can phosphorylate and activate MAPK8/JNK, MAP2K7/MKK7, which suggests a role in the JNK signaling pathway. [provided by RefSeq, Jul 2008], catalytic activity: ATP + a protein = ADP + a phosphoprotein., cofactor: Magnesium., enzyme regulation: Activated by autophosphorylation and homodimerization., function: Activates the JUN N-terminal pathway through activation of the MAP kinase kinase MAP2K7. Acts synergistically with PRDX3 to regulate the activation of NF-kappa-B in the cytosol. This activation is kinase-dependent and involves activating the IKK complex, the IKBKB-containing complex that phosphorylates inhibitors of NF-kappa-B., PTM: Autophosphorylated on serine and threonine residues., sequence caution: Translated as Tyr., sequence caution: Wrong choice of CDS., similarity: Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily., similarity: Contains 1 protein kinase domain., subunit: Homodimer; forms dimers through the leucine-zipper motif. Interacts with the C-terminus of MAPK8IP1 through the kinase catalytic domain. Binds PRDX3. Associates with the IKK complex through the kinase domain., tissue specificity: Expressed in the adult brain, liver, placenta and pancreas, with expression strongest in the pancreas.,