

<b>Product name:</b>	MADCAM1 Rabbit Monoclonal Antibody
<b>Cat number:</b>	MABN84407
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A synthesized peptide derived from human MADCAM1
<b>Reactivity:</b>	Human
<b>Applications:</b>	WB 1:1000-1:2000
<b>Molecular Weight:</b>	Calculated MW: 40 kDa ; Observed MW: 60 kDa
<b>Purification:</b>	Affinity Purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Purified antibody in PBS with 0.05% sodium azide, 0.05% BSA and 50% glycerol.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	Cell adhesion leukocyte receptor expressed by mucosal venules, helps to direct lymphocyte traffic into mucosal tissues including the Peyer patches and the intestinal lamina propria. It can bind both integrin alpha-4/beta-7 and L-selectin, regulating both the passage and retention of leukocytes. Isoform 2, lacking the mucin-like domain, may be specialized in supporting integrin alpha-4/beta-7-dependent adhesion strengthening, independent of L-selectin binding.