
| | |
|--------------------------|---|
| Product name: | MET Mouse Monoclonal Antibody |
| Cat number: | MABN81755 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Monoclonal |
| Concentration: | 1mg/ml |
| Host: | Mouse |
| Isotype: | Mouse IgG1 |
| Immunogen: | Purified recombinant fragment of human MET (AA: 743-932) expressed in E. Coli. |
| Reactivity: | Human,Monkey |
| Applications: | WB 1:500-1:2000,IHC 1:200-1:1000,ICC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400 |
| Molecular Weight: | 155kDa |
| Purification: | Affinity Purification |
| Form: | Liquid |
| Buffer: | Purified antibody in PBS with 0.05% sodium azide |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |
| Background: | <p>This gene encodes a member of the receptor tyrosine kinase family of proteins and the product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival, embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are also associated with multiple human cancers.</p> |