

---

<b>Product name:</b>	MLK1 Rabbit Polyclonal Antibody
<b>Cat number:</b>	ABN13951
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	The antiserum was produced against synthesized peptide derived from human MAP3K9. AA range:561-610
<b>Reactivity:</b>	Human,Mouse
<b>Applications:</b>	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
<b>Purification:</b>	Affinity purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	<p>MAP3K9 (Mitogen-Activated Protein Kinase Kinase Kinase 9) is a Protein Coding gene. Diseases associated with MAP3K9 include retroperitoneal neuroblastoma. Among its related pathways are MAP Kinase Signaling and TGF-Beta Pathway. GO annotations related to this gene include protein homodimerization activity and protein kinase activity. An important paralog of this gene is KSR1.</p> <p>catalytic activity: ATP + a protein = ADP + a phosphoprotein., cofactor: Magnesium., enzyme regulation: Homodimerization via the leucine zipper domains is required for autophosphorylation and subsequent activation., function: Activates the JUN N-terminal pathway., PTM: Autophosphorylation on serine and threonine residues within the activation loop plays a role in enzyme activation. Thr-312 is likely to be the main autophosphorylation site., similarity: Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 1 SH3 domain., subunit: Homodimer., tissue specificity: Expressed in epithelial tumor cell lines of colonic, breast and esophageal origin.,</p>