

---

<b>Product name:</b>	NAA10 Mouse Monoclonal Antibody
<b>Cat number:</b>	MABN81831
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Mouse
<b>Isotype:</b>	Mouse IgG1
<b>Immunogen:</b>	Purified recombinant fragment of human NAA10 (AA: 111-235) expressed in E. Coli.
<b>Reactivity:</b>	Human,Mouse,Monkey
<b>Applications:</b>	WB 1:500-1:2000,ICC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400
<b>Molecular Weight:</b>	26.5kDa
<b>Purification:</b>	Affinity Purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Purified antibody in PBS with 0.05% sodium azide
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	N-alpha-acetylation is among the most common post-translational protein modifications in eukaryotic cells. This process involves the transfer of an acetyl group from acetyl-coenzyme A to the alpha-amino group on a nascent polypeptide and is essential for normal cell function. This gene encodes an N-terminal acetyltransferase that functions as the catalytic subunit of the major amino-terminal acetyltransferase A complex. Mutations in this gene are the cause of Ogden syndrome. Alternate splicing results in multiple transcript variants.