


---

<b>Product name:</b>	NOS2(I/NOS)
<b>Cat number:</b>	AB-90171
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100 ug
<b>Clone:</b>	POLY
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A synthetic peptide corresponding to a sequence at the C-terminus of human nNOS(neuronal)(1418-1434aa IAFIEESKKDTDEVFSS), identical to the related mouse and rat sequences.
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	Western blot: 1:1000-1:2000 Immunohistochemistry (Paraffin-embedded Section): 1:100-1:200
<b>Purification:</b>	Aff. Pur.
<b>Form:</b>	Liquid
<b>Buffer:</b>	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg Thimerosal, 0.05mg NaN <sub>3</sub> .
<b>Storage:</b>	At -20°C for one year.. Avoid repeated freezing and thawing.
<b>Background:</b>	Nitric oxide synthase 1 (neuronal), also known as NOS1, is an enzyme that in humans is encoded by the NOS1 gene. The protein encoded by this gene belongs to the family of nitric oxide synthases, which synthesize nitric oxide from L-arginine. Nitric oxide is a reactive free radical, which acts as a biologic mediator in several processes, including neurotransmission, and antimicrobial and antitumoral activities. In the brain and peripheral nervous system, nitric oxide displays many properties of a neurotransmitter, and has been implicated in neurotoxicity associated with stroke and neurodegenerative diseases, neural regulation of smooth muscle, including peristalsis, and penile erection. This protein is ubiquitously expressed, with high level of expression in skeletal muscle. Multiple transcript variants that differ in the 5' UTR have been described for this gene but the full-length nature of these transcripts is not known. Additionally, alternatively spliced transcript variants encoding different isoforms (some testis-specific) have been found for this gene.



Anti-nNOS(neuronal) antibody,  
Western blotting All lanes: Anti  
nNOS(neuronal) at 0.5ug/ml Lane 1:  
Rat Brain Tissue Lysate at 50ug Lane  
2: MCF-7 Whole Cell Lysate at 40ug  
Predicted bind size: 160KD Observed  
bind size: 160KD



Anti-nNOS(neuronal) antibody, IHC(P) IHC(P): Rat Brain Tissue