

<b>Product name:</b>	NRCAM Mouse Monoclonal Antibody
<b>Cat number:</b>	MABN81211
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Mouse
<b>Isotype:</b>	Mouse IgG1
<b>Immunogen:</b>	Purified recombinant fragment of human NRCAM (AA: 1192-1255) expressed in E. Coli.
<b>Reactivity:</b>	Human
<b>Applications:</b>	ICC 1:20-1:50,ELISA 1:5000-1:20000
<b>Molecular Weight:</b>	144kDa
<b>Purification:</b>	Affinity Purification
<b>Form:</b>	Liquid
<b>Buffer:</b>	Purified antibody in PBS with 0.05% sodium azide
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
<b>Background:</b>	Cell adhesion molecules (CAMs) are members of the immunoglobulin superfamily. This gene encodes a neuronal cell adhesion molecule with multiple immunoglobulin-like C2-type domains and fibronectin type-III domains. This ankyrin-binding protein is involved in neuron-neuron adhesion and promotes directional signaling during axonal cone growth. This gene is also expressed in non-neural tissues and may play a general role in cell-cell communication via signaling from its intracellular domain to the actin cytoskeleton during directional cell migration. Allelic variants of this gene have been associated with autism and addiction vulnerability. Alternative splicing results in multiple transcript variants encoding different isoforms.