

Product name:	NT5C3 Rabbit Polyclonal Antibody
Cat number:	ABN14920
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human NT5C3. AA range:11-60
Reactivity:	Human,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:20000-1:40000
Molecular Weight:	38kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

5'-nucleotidase, cytosolic IIIA(NT5C3A) Homo sapiens This gene encodes a member of the 5'-nucleotidase family of enzymes that catalyze the dephosphorylation of nucleoside 5'-monophosphates. The encoded protein is the type 1 isozyme of pyrimidine 5' nucleotidase and catalyzes the dephosphorylation of pyrimidine 5' monophosphates. Mutations in this gene are a cause of hemolytic anemia due to uridine 5-prime monophosphate hydrolase deficiency. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and pseudogenes of this gene are located on the long arm of chromosomes 3 and 4. [provided by RefSeq, Mar 2012],catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,disease:Defects in NT5C3 are the cause of P5N deficiency [MIM:266120]; also called hemolytic anemia due to P5N deficiency or hemolytic anemia due to UMPH1 deficiency. P5N deficiency is an autosomal recessive condition causing hemolytic anemia characterized by marked basophilic stippling and the accumulation of high concentrations of pyrimidine nucleotides within the erythrocyte. It is implicated in the anemia of lead poisoning and is possibly associated with learning difficulties.,function:Can act both as nucleotidase and as phosphotransferase.,induction:Isoform 2 is induced by interferon alpha in Raji cells in association with lupus inclusions.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,subunit:Monomer.,tissue specificity:Isoform 1 and isoform 3 are expressed in reticulocytes and lymphocytes. Isoform 4 is expressed only in reticulocytes.,