


Product name:	OPA1 (8F7) Rabbit Monoclonal Antibody
Cat number:	MABN15355
Conjugate:	Unconjugated
Size:	100µL
Clone:	8F7
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human OPA1
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:1000-1:5000,IHC 1:200-1:500,IF-P 1:200-1:500
Molecular Weight:	112kDa
Purification:	Affinity purification
Form:	liquid
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Background:

Dynamin-related GTPase required for mitochondrial fusion and regulation of apoptosis. May form a diffusion barrier for proteins stored in mitochondrial cristae. Proteolytic processing in response to intrinsic apoptotic signals may lead to disassembly of OPA1 oligomers and release of the caspase activator cytochrome C (CYCS) into the mitochondrial intermembrane space. Dynamin-related GTPase that is essential for normal mitochondrial morphology by regulating the equilibrium between mitochondrial fusion and mitochondrial fission (PubMed:16778770, PubMed:17709429, PubMed:20185555, PubMed:24616225, PubMed:28746876). Coexpression of isoform 1 with shorter alternative products is required for optimal activity in promoting mitochondrial fusion (PubMed:17709429). Binds lipid membranes enriched in negatively charged phospholipids, such as cardiolipin, and promotes membrane tubulation (PubMed:20185555). The intrinsic GTPase activity is low, and is strongly increased by interaction with lipid membranes (PubMed:20185555). Plays a role in remodeling cristae and the release of cytochrome c during apoptosis (By similarity). Proteolytic processing in response to intrinsic apoptotic signals may lead to disassembly of OPA1 oligomers and release of the caspase activator cytochrome C (CYCS) into the mitochondrial intermembrane space (By similarity). Plays a role in mitochondrial genome maintenance (PubMed:20974897, PubMed:18158317).



Western blot analysis of OPA1 expression in HeLa cell lysate.