

---

|                          |   |
|--------------------------|---|
| <b>Product name:</b>     | p300 Rabbit Polyclonal Antibody   |
| <b>Cat number:</b>       | ABN03402  |
| <b>Conjugate:</b>        | Unconjugated  |
| <b>Size:</b>             | 100µL   |
| <b>Clone:</b>            | Polyclonal  |
| <b>Concentration:</b>    | 1mg/ml  |
| <b>Host:</b>             | Rabbit  |
| <b>Isotype:</b>          | IgG   |
| <b>Immunogen:</b>        | Synthesized peptide derived from part region of human protein. AA range 1520-1570   |
| <b>Reactivity:</b>       | Human,Mouse,Rat   |
| <b>Applications:</b>     | WB 1:500-1:1000,IHC 1:50-1:100,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000   |
| <b>Molecular Weight:</b> | Calculated MW: 264 kDa; Observed MW: 300 kDa  |
| <b>Purification:</b>     | Affinity Purification   |
| <b>Form:</b>             | Liquid  |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |
| <b>Background:</b>       | EP300: E1A binding protein p300. This gene encodes the adenovirus E1A-associated cellular p300 transcriptional co-activator protein. It functions as histone acetyltransferase that regulates transcription via chromatin remodeling and is important in the processes of cell proliferation and differentiation. It mediates cAMP-gene regulation by binding specifically to phosphorylated CREB protein. This gene has also been identified as a co-activator of HIF1A (hypoxia-inducible factor 1 alpha), and thus plays a role in the stimulation of hypoxia-induced genes such as VEGF. Defects in this gene are a cause of Rubinstein-Taybi syndrome and may also play a role in epithelial cancer. |