

---

|                          |  |
|--------------------------|--|
| <b>Product name:</b>     | PARP Mouse Monoclonal Antibody   |
| <b>Cat number:</b>       | MABN80824  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Mouse  |
| <b>Isotype:</b>          | Mouse IgG1   |
| <b>Immunogen:</b>        | Synthetic peptide of human PARP, conjugated to KLH.  |
| <b>Reactivity:</b>       | Human  |
| <b>Applications:</b>     | WB 1:500-1:2000,ELISA 1:5000-1:20000,FC 1:200-1:400  |
| <b>Molecular Weight:</b> | 117kDa   |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide.  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |
| <b>Background:</b>       | <p>This gene encodes a chromatin-associated enzyme, poly(ADP-ribose)transferase, which modifies various nuclear proteins by poly(ADP-ribose)ation. The modification is dependent on DNA and is involved in the regulation of various important cellular processes such as differentiation, proliferation, and tumor transformation and also in the regulation of the molecular events involved in the recovery of cell from DNA damage. In addition, this enzyme may be the site of mutation in Fanconi anemia, and may participate in the pathophysiology of type I diabetes.</p> |